

# **ACE Inhibitors**

<b>Medication Name</b>		<b>Brand Name</b>				
Captopril	0	Acenorm	0	Captopril	0	Topace
	0	Capoten	0	Captohexal	0	Zedace
Enalapril	0	Acetec	0	Auspril	0	Enalapril
	0	Alphapril	0	Enahexal	0	Enalapril
	0	Amprace	0	Enalabell		Maleate
					0	Renitec
Fosinopril	0	Fosipril	0	Fosinopril	0	Monace
	0	Fosinopril		Sodium	0	Monopril
Lisinopril	0	Fibsol	0	Lisinopril	0	Lisodur
	0	Liprace	0	Lisinopril	0	Lisoril
	0	Linopril		hexal	0	Prinivil
	0	Lisinobell	0	Lisinotrust	0	Zinopril
			0	Lisodrate	0	Zestril
Perindopril	0	Coversyl	0	Ozapace	0	Perindopril
	0	Indaprex	0	Perindo		Arginine
	0	Indopril	0	Perindobell	0	Perindopril
	0	Indosyl	0	Prexum		Erbumine
		Mono				
Quinapril	0	Accupril	0	Aquinafil	0	Filpril
	0	Acquin	0	Asig	0	Qpril
Ramipril	0	Prilace	0	Ramibell	0	Tryzan
	0	Ramace	0	Titrace	0	Vascalace
Trandolapril	0	Dolapril	0	Goricap	0	Tranalpha
	0	Gopten	0	Odrik	0	Trandolapril

#### Darwin

Shop 1 & 2 Tiwi Place, Tiwi NT 0810 PO Box 40113, Casuarina NT 0811 Phone: 08 8927 8488 Fax: 08 8927 8515 E: info@ healthylivingnt.org.au

### Alice Springs

Jock Nelson Centre, 7/16 Hartley Street, Alice Springs NT 0870 Phone: 08 8952 8000 Fax: 08 8952 7000 E: alicesprings@ healthylivingnt.org.au

www.healthylivingnt.org.au ABN 11 374 693 055

Healthy Living NT is the trading name of the Diabetes Association of the Northern Territory Incorporated.

### What is it used for?

ACE Inhibitors are used to lower blood pressure and improve heart failure. They help your heart by relaxing blood vessels, therefore making it easier for your heart to pump blood around the body.

## How to take this medication

This medication must be taken regularly. If a dose is missed, only take the dose if you remember within three hours. If not, take the next dose when it is due normally. Do not take the missed dose or two doses at once.

Take this medication only as the directions say on the pharmacy label.

## Before taking this medication it is important to tell your Doctor if you:

- Are pregnant or plan to become pregnant
- Are breast feeding
- Have had previous reactions to other ACE Inhibitors
- Have kidney disease

### **Side Effects**

Side effects occur in some people taking this medication. If they are troublesome to you, speak to your doctor or pharmacist.

- Dizziness, light-headedness
- Loss of taste
- Headache
- Persistent cough
- Unusually tired
- Diarrhoea

## Other important facts

- Dizziness/ light-headedness may occur if you get up suddenly. This is due to the effect of the medication on your blood pressure
- Always make sure you get up slowly from a lying or sitting position
- Only take potassium supplements (Slow K®, KSR®) if your doctor has told you to
- Tell your doctor all the medications you take, including those you get with out a prescription
- Never share your medicines with anyone
- Keep the medicine out of reach of children

If you have any queries about your medications, talk to your Doctor or pharmacist.

This information is only designed for use as an educational aid and is general in nature.

This resource was produced using information from Hospital Pharmacy Services and current medication guidelines.