

Your partner in heart & diabetes health

healthylivingNT

Beta Blockers

Medication name		Brand Name				
Atenolol	0	Anselol	0	Atenolol Gen Rx	0	Tenolten
	0	Atehexal	0	Colaptyn	0	Tenormin
	0	Aten	0	Noten	0	Tensig
Bisoprolol	0	Beprol	0	Biso	0	Bisoprolol Gen
	0	Bicard	0	Bisoprolol		Rx
	0	Bicor		Fumarate	0	Bispro
Carvedilol	0	Carvedilol	0	Dicarz	0	Kredex
	0	Carvedilol	0	Dilasig	0	Vedilol
		Gen Rx	0	Dilatrend	0	Volirop
Labetolol	0	Presolol	0	Trandate		
Metoprolol	0	Betaloc	0	Metolol	0	Metrol XL
	0	Lopresor	0	Metoprolol	0	Minax
	0	Metatar	0	Metoprolol BC	0	Minax XL
	0	Metohexal	0	Metrol	0	Toprol XL
Nebivolol	0	Lobivon	0	Nebilet	0	Nebivolol
Propranalol	0	Deralin	0	Hemangiol	0	Propranalol Gen
	0	Drofate	0	Inderal		Rx
Sotalol	0	Cardol	0	Sotab	0	Sotalol Gen Rx
	0	Sotacor	0	Sotacor	0	Sotalol
	0	Solavert	0	Sotahexal		Hydrochloride

What is it used for?

Beta blockers are used to treat high blood pressure, irregular heart rhythms, help prevent angina and to reduce complications after a heart attack. Beta blockers open up blood vessels, which reduces blood pressure, and prevents some nerve impulses from speeding up your heart beat. This also prevents the heart from working too hard.

Darwin

Shop 1 & 2 Tiwi Place, Tiwi NT 0810 PO Box 40113, Casuarina NT 0811 Phone: 08 8927 8488 Fax: 08 8927 8515 E: info@ healthylivingnt.org.au

Alice Springs

Jock Nelson Centre, 7/16 Hartley Street, Alice Springs NT 0870 Phone: 08 8952 8000 Fax: 08 8952 7000 E: alicesprings@ healthylivingnt.org.au

www.healthylivingnt.org.au ABN 11 374 693 055

Healthy Living NT is the trading name of the Diabetes Association of the Northern Territory Incorporated.

How to take this drug

This medication must be taken regularly. If a dose is missed, only take the dose if you remember within three hours. If not, do not take the missed dose, and take the next dose when it is due normally. Do not take two doses at once.

Take this medication only as the directions say on the pharmacy label.

Before taking this medication it is important to tell your doctor if you:

- Are pregnant or plan to become pregnant, are breast feeding
- Have had previous allergic reactions to other beta blockers
- Have asthma, or other difficulties breathing
- Have low blood pressure
- Have muscle weakness
 - Have kidney, liver disease or diabetes

Side Effects

Side effects occur in some people taking this medication. If they are troublesome to you, speak to your doctor or pharmacist.

- Drowsiness/tiredness, trembling/shaking
- Headache, aches/pains, flushes
- Difficulty sleeping, confusion
- Mood changes, depression
- Loss of appetite/nausea, weight gain, stomach upset, diarrhoea, constipation
- Problems with sexual function

Other important factors

- Dizziness/light-headedness may occur if you get up suddenly. This is due to the effect of the drugs on your blood pressure.
- Always make sure you get up slowly from a lying or sitting position
- Your doctor or dentist needs to know you are on this medication before any procedures or operations
- Tell your doctor all the medications you take including those you get without prescriptions
- Never share your medications with anyone
- Keep the medicine out of reach of children

If you have any queries about your medications, talk to your Doctor or pharmacist. This information is only designed for use as an educational aid and is general in nature.

This resource was produced using information from Consumer Medical Information and assistance from Royal Darwin Hospital Pharmacy.